WISELEAF SCIENTIFIC VENTURES

COMPREHENSIVE PREGNANCY CARE: AN EVIDENCE-BASED MULTIDISCIPLINARY APPROACH

Chapter 3: Nutrition and Lifestyle Management during Pregnancy: An Evidence-Based Approach

Dr. Richa Srivastava

Professor, Department of Pharmaceutics, Ambekeshwar Group Of Institutions, Kathwara, Maa Chandrika Devi Road, Bakshi Talab, Lucknow, UP, India

Mail id - lilllyricha@gmail.com

Abstract

The gestational period is a vital time that comes with increasing nutritional, physical, and psychological demands that impact mother and baby. Good nutrition and healthy lifestyle choices are crucial to ensure proper functioning of the placenta, good fetal development, appropriate weight gain during pregnancy and successful outcomes. The present chapter provides an evidence-based review of nutrition and lifestyle management during pregnancy focusing on the dietary needs in each trimester and practical advice on how to stay healthy. The author elaborates on the energy needs and the functions of macronutrients: carbohydrates, fats, and proteins, as well as micronutrients such as zinc, calcium, vitamin D, vitamin B12, iodine and iron. The chapter also discusses how to avoid nutritional deficiencies by having a diversified diet, taking supplements and regular diet assessment throughout pregnancy. The section continues by examining how regular physical activity, yoga before childbirth, meditation, and relaxation skills can improve physical and mental health during pregnancy. Important issues related to lifestyle such as healthy weight gain during pregnancy, quitting smoking and drinking alcohol, consumption of coffee, food safety, sufficient sleep, drinking enough water, and taking medicine and dietary supplements safely are studied.

Keywords: Pregnancy; Maternal Nutrition; Nutritional Requirements; Physical Activity; Maternal Well-being.

3.1 Introduction

Pregnancy refers to a different physiological state in which a range of anatomical, hormonal, metabolic, and psychological changes occur in the body that leads to the fetus's development and prepares the mother for delivery and lactation. Changes in physiology increase nutritional and lifestyle needs for future mothers. Hence, good nutrition and correct behaviours are

important constituents of prenatal care. Maternal nutritional condition at the moment of conception and subsequently during pregnancy is critical since it impacts maternal health, functioning of the placenta, development of the fetus, and pregnancy outcomes. Good nutrition is necessary not only for meeting the needs of the pregnant woman but also for ensuring the child's health in the long term. During gestation, the energy and nutrient requirements of the mother and fetus keep changing according to gestational age. For many functions like tissue growth, erythropoiesis, bone development, brain development and metabolism several nutrients, among which are protein, iron, folate, calcium, vitamin D, iodine, vitamin B12, zinc and other microelements, become necessary. [1] A too poor or uneven diet can cause nutritional deficiency and lead to maternal anemia, lack of weight gain, health problems in the mother, fetal growth restriction, premature delivery and low birth weight. Conversely, excessive energy intake during pregnancy could lead to development of gestational diabetes and hypertensive issues as well as hyperglycemia, Cesarean section and retention of excessive weight postdelivery. Therefore, reaching the balance between sufficient nutrient intake and appropriate weight gain is both desirable and beneficial during pregnancy. [2]

Dietary quality is important because the desired increase in nutritional needs should be satisfied by a wide and nutrient-rich dietary intake. A consistent intake of whole grains, vegetables, fruits, legumes, dairy products, eggs, low-fat protein sources, nuts, and seeds can guarantee the intake of a wide variety of nutrients. There are cases in which nutritional supplementation should be required, i.e., when dietary intake is inadequate for any reason or when specific insufficiency or nutritional risk is revealed. Another factor to keep in mind in nutritional counseling is individual specifics of clients which can include pre-pregnancy BMI, age, food habits, food availability, and the presence of any medical conditions. In addition to sufficient nutrition during pregnancy, women with no contraindications should practice regular exercise. Adequate physical activity helps achieve optimal fitness status, ensures healthy weight during pregnancy, promotes healthy cardiovascular conditions, and provides psychological well-being. Appropriate exercises may include, but are not limit to, walking, swimming, stationary cycling, and pelvic floor exercises instead of any physical activity that is considered moderate for pregnancy. It is advisable to alter the exercise program as pregnancy advances, increasing hydration rate and careful observation of environmental conditions, overall physical comfort, as well as possible warning signals for resuming physical activity and seeking medical advice. [3]

The period when a woman goes through pregnancy is significant with respect to the provision of mental and emotional well-being. Hormonal changes, bodily changes, discomfort, worries about the birth, family duties, financial conditions, and acclimatization to being a parent influence the mental state. Stress and anxiety and sleep disturbances affect the quality of life and interfere with normal activities. Techniques such as mindfulness, meditation, relaxation,

sleep, help from people surrounding a person, etc. can help improve mental well-being. It is vital to detect psychological disorders as soon as possible so that a woman can get professional assistance promptly. Insisting on a healthy lifestyle involves more than just a balanced diet and physical activity; it also means giving up all tobacco, nicotine, alcohol, and any harmful substance. It is essential to control caffeine intake, maintain proper hydration levels, ensure good food and drink hygiene, prepare and store food in a safe manner, enjoy sufficient sleep, and practice responsible medicine and dietary supplements consumption. Women in a delicate condition should receive advice concerning medication and supplements for problems they may face, because some substances may have adverse effects on the fetus. [4]

Healthcare professionals play a key role in incorporating the proposed recommendations into routine antenatal care practice.[5] Obstetricians, doctors, nurses, midwives, nutritionists, pharmacists, and other healthcare professionals can assess nutrition status, track weight gain during pregnancy, and contact mother with appropriate diet and lifestyle tips. Regular antenatal checks give an opportunity to promote healthy habits, evaluate factors preventing good nutrition and physical activity, and change recommendations according to what is needed by the mother and fetus. Thus, it is essential for the selected model of pregnancy management to adopt a comprehensive approach, which takes into account proper nutrition, physical activity, mental well-being, sufficient rest, and healthy lifestyle. This comprehensive model can help to prevent nutritional deficiencies, as well as promote proper weight gain during pregnancy, minimize alterable risk factors, and improve maternal and fetal outcomes. This chapter provides an evidence-based insight into the nutritional needs during pregnancy, trimester-based diet planning, vitamins and minerals needed during pregnancy, prevention of inadequate nutrition, exercises, yoga and meditation, mental well-being, and modifications in lifestyle. In addition, importance of individualized counselling and constant monitoring of health is highlighted.

3.2 Nutritional Requirements during Pregnancy

During pregnancy, the body's nutritional demands heightened to ensure maternal bodily changes, placenta formation, fetal development, and readiness for lactation. A proper diet containing appropriate amount of energy, macronutrients, micronutrients, and fluid is important to ensure the well-being of both the mother and baby.

3.2.1 Energy Requirements

As mentioned in (Table 3.1) as the pregnancy moves ahead, the energy demand increases. There is no particular need to increase energy intake in the first trimester, whereas average energy intake may need to be increased by about 340 kcal/day in the second trimester and 450 kcal/day in the third trimester. It is best if the energy comes from nutrient-rich foods, such as whole grain, fruits, vegetables, legumes, dairy products, nuts, and lean sources of protein. Both

insufficient and excessive energy intake should be avoided for the right weight gain during pregnancy.[6]

Table 3.1: Energy and Nutritional Requirements during Pregnancy

Nutritional Component	First Trimester	Second Trimester	Third Trimester	Major Role/Consideration
Energy	No substantial additional energy required	**+340 kcal/day**	**+450 kcal/day**	Supports fetal growth, maternal tissue expansion and preparation for lactation
Carbohydrates	~175 g/day	~175 g/day	~175 g/day	Primary energy source; supports fetal growth and protein sparing
Protein	≥1.1 g/kg/day	≥1.1 g/kg/day	≥1.1 g/kg/day	Supports fetal, placental and maternal tissue growth
Fats	Adequate intake; emphasize unsaturated fats	Emphasize unsaturated fats and omega-3 fatty acids	Emphasize unsaturated fats and omega-3 fatty acids	Provides energy and supports fetal brain and visual development
Micronutrients	Increased focus on folate, iron, iodine, B12 and vitamin D	Increased focus on protein, iron, calcium, folate, vitamin D, iodine and omega-3	Continued emphasis on iron, calcium, vitamin D, iodine and omega-3	Supports maternal health, placental function and fetal development
Fluids	Adequate hydration	Adequate hydration	Adequate hydration	Supports blood volume, amniotic fluid, nutrient transport and temperature regulation

3.2.2 Macronutrients

Carbohydrates

The biggest source of energy is carbohydrates which also help save proteins for developing tissue. A recommendation for carbohydrates consumption during pregnancy is about 175 g/day. It is better to eat whole grain products, fruits and vegetables, and legumes. Foods with additional sugar must be consumed as low as possible. [7]

Proteins

Protein is very important for fetal growth, and for the development of placenta and maternal tissues. The suggested amount of protein intake is not less than 1.1 g/kg of weight/day in pregnant women. Some of the sources of protein are pulses, beans, eggs, dairy, fish and poultry, and nuts.

Fats

Dietary fats are responsible for giving energy as well as for the development of brain and eyesight. Unsaturated fats and omega-3 fatty acids, particularly DHA, are necessary and are consumed through different foods including nuts, seeds, vegetable oils, avocados, and low mercury fish. It's important to reduce foods with saturated and trans fats. [8]

3.2.3 Micronutrients

Micronutrients are crucial to fetal development and supporting the health of the mother and metabolism. Vitamins that are particularly important include folate, vitamins A and B12, as well as vitamins C, D, B6, and E. The most important minerals include iron, calcium, iodine, zinc, and magnesium. A balanced diet of green vegetables, fruits, whole grains, legumes, dairy products, nuts, seeds, eggs, and relevant animal products can help to cover these micronutrient needs. Supplements may be necessary if dietary habits lead to a deficiency. [9]

3.2.4 Water and Hydration

Proper hydration aids in enhancing blood volume, generating amniotic fluid, transporting nutrients, digesting food, and regulating body heat. An average amount of water consumed around 3.0 L/day is recommended during pregnancy, however, this varies from person to person. Water should always be the preferred method of hydration and consumption of sugary drinks and alcohol should be avoided while caffeine must be limited as per recommendations. [10] Overall, balanced energy and macronutrients along with other essential micro nutrients and adequate hydration is necessary for healthy nutrition at the time of pregnancy. [11]

3.3 Month-wise Diet Planning During Pregnancy

Dietary requirements gradually modify during pregnancy because of the needs of the growing fetus, developing placenta, expanding maternal tissues, and preparation for lactation. As shown in (Fig 3.1) it is important that all aspects of nutrient intake energy, protein, fatty acids, vitamins, minerals, fibre, and fluid—are sufficient in a balanced diet. A pregnant woman should carefully plan her daily meals based on food variety and nutrient density while taking into account her pre-pregnancy eating habits and dietary preferences, her culture, and her state of health. [12]

3.3.1 First Trimester (Months 1–3)

Pregnancy's first trimester is marked by several changes in the body, including nausea, vomiting, tiredness, and appetite alterations. This period in a woman's life, however, does not require many extra calories, but rather an adequate intake of specific nutrients, including folate, iron, iodine, vitamin B12, and vitamin D. It is advisable to consume whole grains, pulses, vegetables, fruits, dairy, eggs, and nuts. If a person is facing nausea due to pregnancy, proper nutrition is important because it helps to cope with the issue.

3.3.2 Second Trimester (Months 4–6)

As pregnancy progresses into the second trimester, fetal growth accelerates and mother's blood volume and tissue requirements increase. Energy requirements, in general, increase by about 340 kcal per day, although it should be noted that individual requirements depend on a person's pre-pregnancy BMI, physical activity level, and weight gain during pregnancy. Pregnant women should make sure their diet is rich in protein, iron, calcium, folate, vitamin D, iodine, and omega-3 fatty acids. In order to meet these nutrient needs on a regular basis, a pregnant woman is expected to eat whole grain products, legumes, vegetables, fruit, dairy products, eggs, lean meat, nuts, seeds, and fish.

3.3.3 Third Trimester (Months 7–9)

The third trimester involves fast growth of the fetus, the enhancement of fetal organs, and growing food requirements of the mother. The caloric needs may be increased by up to 450 kcal a day, depending on the situation. The pregnant woman needs to take in enough protein, iron, calcium, vitamin D, iodine, and omega-3 fatty acids. A pregnant woman should eat wholemeal cereals, vegetables, fruits, grains, dairy products, eggs, low-fat meats and fish. For women suffering from heartburn, regurgitation, stomach pain, or low appetite in late pregnancy, smaller and more frequent meals are suggested..

3.3.4 Foods to Include

Nutrient-packed food from different groups should be included in healthy pregnancy diet. Whole grains and millets are good sources of carbohydrates and fibre. Protein-rich pulses, beans, lentils, eggs, dairy products, lean meats, poultry, and low mercury fishes help in getting the other important nutrients. Different colored fruits and vegetables are the good sources of vitamins, minerals, antioxidants, and fibres. Milk, curd, cheese, and other enriched substitutes help in getting calcium and other nutrients. Nuts, seeds, avocadoes, and healthy vegetable oils provide unsaturated fats and omega 3 fatty acids. During the whole pregnancy, one should drink sufficient amounts of water. [13]

3.3.5 Foods to Limit or Avoid

Pregnancy requires avoidance or limitation of certain foods and drinks to lessen the risk of getting an infection as well as harm to maternal and fetal health. Alcohol should always be avoided during pregnancy.[14] In addition, raw or undercooked meat, poultry, eggs and fish should be avoided as this could lead to foodborne infection. Fish containing high levels of mercury should not be consumed; in contrast, fish with low mercury levels can be consumed as recommended. Moreover, pregnant women should limit caffeine intake and minimize highly processed food, as this food can lead to excessive intake of sugar, sodium, saturated and trans fats. Finally, fruits and vegetables should be cleaned carefully before being eaten and proper food storage and preparation techniques should be used to minimize the risk of foodborne illnesses. [15]




Fig 3.1: Trimester-Wise Dietary Priorities during Pregnancy

3.4 Role of Essential Nutrients

Proper nutrition, which is crucial during pregnancy, is required not only for the health of the mother, but also for the growing fetus. While pregnant, women undergo the increase of blood volume, development of the placenta and the fetal organism and preparation for breastfeeding. As shown in (Fig 3.2) Therefore, it is necessary to consume well-balanced diet during pregnancy so that these special needs may be satisfied.

3.4.1 Iron

During pregnancy, iron plays an important role in the generation of hemoglobin and the transport of oxygen to both maternal and fetal tissues. The needs for iron increase substantially due to the increase in blood volume required by the mother, and the production of red blood cells in fetal life. Adequate intake of iron will avoid iron deficiency anemia and its consequences such as fatigue of the mother, decreased ability to work, preterm delivery or low birth weight of the baby. Foods that contain iron are dark green leafy vegetables, legumes, beans, lentils, fortified cereals, eggs, lean meat and poultry. Vitamin C intake facilitates the absorption of iron obtained from plant resources as well. [16]

3.4.2 Calcium

Calcium plays an important role in the formation of teeth and bones in the fetus and supports the proper functioning of the nervous system and heart. In addition, pregnant women need an adequate supply of calcium, not only to satisfy the increased needs of the fetus but also to maintain healthy bones. Dairy products such as milk, curd/yogurt, and cheese, as well as certain plant drinks, sesame seeds, almonds, and some leafy greens are great sources of calcium. Adequate intake of calcium can also prevent hypertensive disorders during pregnancy in women with low intake of this mineral. [17]

3.4.3 Folic Acid

Folate is a crucial vitamin that helps in DNA formation, cell development, and formation of the placenta and the child. Proper folate intake before pregnancy is key to prevent neural tube problems. Folate can be obtained via consumption of spinach, green salad, citrus fruits, nuts, and fortified grains. Because of the increased need for folic acid during pregnancy, women are recommended to take pills with folic acid before getting pregnant. Women with a higher risk of neural tube problems are advised to take pills before pregnancy as well according to recommendations. [18]

3.4.4 Vitamin D and Other Vitamins

Vitamin D plays a crucial role in metabolism of the calcium and phosphate, formation of bones and functioning of the immune system. Sufficient vitamin D levels in mothers help the fetus in skeleton development and mothers in protecting the health of their bones. The most important sources of vitamin D are sunlight exposure, fortified food, eggs, and some fatty fish, with supplementation recommended for women with lower intake of vitamin D or its deficiency. Other vitamins are also crucial regarding pregnancy. Vitamin A helps vision, immune system, and cell development; vitamin B12 is critical for DNA synthesis and neurological development; vitamin C is responsible for collagen production, immune system, and absorbing iron; and vitamin B6 plays a role in protein metabolism and helps the nervous system.

3.4.5 Other Essential Micronutrients

Numerous additional micronutrients are essential to the good development of the mother and child. Iodine contributes to the manufacture of thyroid hormones and is especially important for the brain of the developing fetus. Zinc promotes the synthesis of DNA, maintains the process of division of cells, and functions of the immune system. Magnesium is involved in many important biochemical processes and is responsible for the normal performance of muscles and nervous system processes. Choline is necessary for the cell membrane formation and brain development, whereas omega-3 fatty acids (in particular DHA) contribute to the development of the brain and vision of the fetus. You can obtain all these substances from the consumption of fish, eggs, dairy, legumes, nuts and whole grains.




Fig 3.2: Key Micronutrients and Their Roles in Maternal and Fetal Health

3.5 Prevention of Nutritional Deficiencies

Nutritional deficiencies frequently occur in pregnancy as energy, protein, vitamins, and minerals are required in larger amount than before. Inadequate dietary intake, restricted food diversity, and still higher dietary requirements may be a risk factor for nutritional deficits. However, proper nutritional evaluation of a diet during the early stages of pregnancy will help avoid deficits and promote maternal and fetal health.

3.5.1 Common Nutritional Deficiencies

As mentioned in (Table 3.2) the most serious and relevant deficiencies are recorded among pregnant women include iron, folate, vitamin B12, vitamin D, calcium, and iodine deficiencies. Iron deficiency is especially widespread and can lead to anaemia, fatigue, decreased activity levels and higher chances of adverse pregnancy outcomes. Folate and vitamin B12 deficiencies influence DNA synthesis and fetal development. Vitamin D and calcium deficiencies can negatively impact the bone health of both mother and fetus. Iodine deficiency may affect production of the thyroid hormones and fetal nervous system development. The risk of deficiencies is higher in women with limited diets, inadequate intake, malabsorption and increased nutritional requirements.

Table 3.2: Common Nutritional Deficiencies during Pregnancy and Their Consequences

Nutrient	Major Causes/Risk Factors	Maternal Consequences	Fetal/Neonatal Consequences
Iron	Inadequate dietary intake; increased pregnancy requirements	Anaemia, fatigue, reduced activity levels	Increased risk of adverse pregnancy outcomes
Folate	Inadequate dietary intake; increased nutritional requirements	Impaired folate status and risk of deficiency	Impaired DNA synthesis and fetal development; increased risk of neural tube defects
Vitamin B12	Limited dietary intake; inadequate nutrient availability	Anaemia and impaired cellular functions	Impaired neurological development and fetal growth
Vitamin D	Inadequate intake or limited nutritional status	Poor bone health	Impaired fetal skeletal development
Calcium	Inadequate dietary intake	Compromised maternal bone health	Impaired fetal bone development
Iodine	Inadequate dietary intake	Impaired thyroid hormone production	Impaired fetal nervous system and brain development

3.5.2 Dietary Strategies

The first way to avoid nutritional deficiencies during pregnancy is to consume a varied and balanced diet course. Meals must include whole grains, pulses and legumes, fruits and vegetables, dairy products or fortified options, eggs, low-fat proteins, nuts, and seeds. Iron-containing food must be paired with vitamin C food in order to enhance iron absorption. Calcium-containing food should be eaten regularly to satisfy the increased needs. Fortified food products may be beneficial when there is not enough food sources of the nutrients.[19]

Pregnant women should be taught proper meal times, food portions, their need for drinking, food safety, and making healthy food choices. [20]

3.5.3 Nutritional Supplementation

The nutrition gained from food may not necessarily satisfy the increasing requirements during the gestational period. Hence, some nutrient supplements including folic acid, iron, calcium, vitamin D, iodine or vitamin B12 may be prescribed depending on the individual needs and local clinical guidelines. The use of folic acid is essential before conception and during early gestation to minimize the chances of neural tube defects, while iron supplementation along with folic acid is commonly used to treat and prevent anemia. The intake of calcium and vitamin D may become necessary when dietary consumption or nutritional status is insufficient.

3.5.4 Monitoring Nutritional Status

Regular assessment of nutrition is useful for detecting nutritional deficiency as well as taking appropriate measures timely. Assessment can include dietary recall, pre-pregnancy BMI, weight gain during pregnancy, physical examination, and relevant laboratory testing. Indicator of level of hemoglobin and other laboratory tests may be used to find anemia or certain deficiency illnesses if needed based on clinical signs and symptoms. Monitoring of maternal weight during pregnancy is important in order to keep it within recommended levels. Nutritional advice should be periodically updated taking into account gestational age changes, dietary patterns, and laboratory results. The success of prevention of nutritional deficiency lies in the complex approach. The issues can be solved by means of proper nutrition and education.

3.6 Exercises during Pregnancy

As shown in (Fig 3.3) Regular physical activity is a crucial part of a healthy pregnancy for women who do not have any medical restrictions. Regular exercise can maintain fitness levels, promote healthy gain in gestational weight and enhance the woman's mood and sleeping patterns. The recommendations concerning the types and intensity of exercises should be personalized based on the woman's activity level prior to pregnancy, the gestational age, health condition and any complications related to pregnancy. [21]

3.6.1 Benefits of Physical Activity

Exercise during pregnancy requires consistency to help improve cardiovascular fitness, muscular strength, flexibility, and overall well-being. It also helps to keep excessive gestational weight in check and decreases the risk of developing several pregnancy-related conditions like gestational diabetes, hypertension, and other pains associated with pregnancy. Furthermore, physical activity positively influences sleep, anxiety, and mental health. A proper routine of exercises prepares the future pain of childbirth and triggers postnatal recovery. [22]

3.6.2 Recommended Exercises

For the majority of healthy pregnant women, a minimum of 150 minutes of moderate-intensity aerobic exercise on a weekly basis is commonly recommended and should be spread over a number of days. Walking, swimming, using a stationary bicycle, doing low-impact aerobics, and participating in modified prenatal exercise programs are just a few viable options. [23] Muscles strengthening activities involving major muscle groups can be performed as well, so long as intensity is appropriate. Exercises focusing on the pelvic floor can be included as well as they help to strengthen pelvic floor muscles and ensure normal urination. [24]

3.6.3 Exercises to Avoid

It is usually advisable for expecting mothers to steer clear of activities that expose them to a greater risk of falling, abdominal injury, or too much physical stress. Such activities may be those conducted in contact sports, activities that pose a major risk of falling, diving under the sea, and engaging in high-intensity activities that exceed one's physical capability. One should also avoid continuing activities in hot and humid conditions to avoid overheating and dehydration. With the progression of pregnancy, activities that involve prolonged lying flat on the back should be adjusted in accordance with either comfort or clinical recommendations. [25]

3.6.4 Safety Precautions

Prenatal exercise must begin gradually and be modified according to the degree of gestation. Sufficient hydration must be ensured, suitable clothing must be worn, comfortable footwear must be used, and the environment must be comfortable. Women must refrain from being exhausted after exercising, and rest must be provided. Medical advice must be sought before doing or continuing any exercise in case of any medical or obstetric issue. Exercise must be stopped and medical advice must be sought in case of warning signs like vaginal bleeding, leakage of amniotic fluid, regular and painful contractions, dizziness and fainting, chest pain, struggling to breathe, severe headache, or any complications in fetal movement. [26]

3.7 Yoga and Meditation during Pregnancy

To have a fit pregnancy, yoga and meditation are to be included as additional techniques if suitable modifications are made for pregnant women and they are practiced under proper supervision. Prenatal yoga is mild exercises, stretching, breathing techniques, and relaxation techniques which can be beneficial for both physical and mental aspects of future mothers. Mindfulness techniques can help pregnant women deal with tension during pregnancy. One should not substitute ordinary reproductive health care and prescribed medications with these practices. [27]



Fig 3.3: Safe Physical Activity During Pregnancy

3.7.1 Benefits of Prenatal Yoga

Prenatal yoga can aid in increasing flexibility, strength, balance, and bodily awareness while alleviating common pregnancy-related issues like labor pain and body strain and increasing fatigue. Yoga can also help with relaxation and sleep improvement while boosting emotional well-being. The various yoga poses practiced may assist women in learning ways to deal with various stressful situations. The extent and kind of yoga should be changed based on the current pregnancy stage and the ability of the individual. [28]

3.7.2 Safe Yoga Practices

It is important for any pregnancy yoga exercise to involve mild movements, active stretching, proper breathing, as well as relaxation rather than demanding positions. Possible examples of this could be standing, sitting, or lying on the side while using adjustments as required. All these exercises have to be carried out slowly and expertly, as pregnancy could lead to increased muscle flexibility due to the system's hormonal changes. With the passage of time, some pillows, blocks, or other supports could be used for better comfort and stability. It is advisable to carry out pregnancy yoga also under a professional instructor's supervision who is trained in pregnancy-related modification options. [29]

3.7.3 Meditation and Relaxation Techniques

Employing the techniques of meditation and mindfulness can prove to terminate the anxiety caused by pregnancy in expectant mothers. It is enough to give a few minutes of our time to using some techniques. These techniques may include breathing exercises, body scan, progressive muscle relaxation, guided imagery and mindfulness meditation, all of which are convenient and easy to bring to life on a regular basis. Breathing deeply and slowly can eliminate body tension and fill the mind with tranquility. Regular relaxation practices along with social support and healthy lifestyle can ensure good sleep and well-being of an individual.

3.7.4 Safety Considerations

While modified yoga can be good for many healthy pregnant women, it is still important for individual practice to take personal situation into account. Poses which may lead to pain, dizziness or breathlessness, and abdominal discomfort should not be performed. Pregnant women should avoid practicing hot yoga, forceful breathing techniques, deep abdominal compression, and any extreme stretch. Prenatal women suffering from some medical complications should consult their doctor before practicing yoga or organized exercise. Women should stop practicing yoga if they experience any warning signs such as vaginal bleeding, leaking, painful contractions, dizziness, chest pain, or strong shortness of breath.

3.8 Mental Well-being during Pregnancy

Mental health is an important part of healthy pregnancy since women go through many changes in their bodies, hormones, emotions, and social life during antenatal period. Pregnancy can evoke positive feelings such as happiness and enjoyment, but it can also cause negative emotions, including worries, doubts, anxiety, and concerns about the pregnancy, childbirth, and motherhood itself, body image, finances, and relationships with family. The psychological well-being of mothers can affect other aspects of pregnancy, including sleeping, eating habits, physical activity, and antenatal care.

3.8.1 Emotional Changes

It is normal for emotional alterations to occur during pregnancy, and they can differ from one individual to another as well as across various stages. Changes in the hormone levels, symptoms of pregnancy, body alterations, worries about the unborn child's health, fear of delivering, as well as the transition into parenthood can have an impact on emotions. Various moods can be observed in ladies, such as irritability, fear, as well as unstable emotions during these periods. The fact that the woman's mood has changed does not necessarily mean that something is wrong; her felling sadness, constant worries, being unable to find pleasure, experiencing difficulties with day-to-day activities can necessitate a period of examination. [30]

3.8.2 Stress and Anxiety Management

Pregnancy-related stress manifests in apprehension over issues related to mother and child's welfare, childbirth, financial problems, family responsibilities, or changes in one's personal or professional life. Minor stress can typically be managed through healthy lifestyles, while severe stress can influence sleep, general life wellness, and ability to function daily. Among efficient strategies are regular reasonable physical activity, special breathing techniques, as well as mindfulness and other relaxation methods. [31]

3.8.3 Sleep and Rest

Adequate sleep and regular times for resting help maintain energy, balance of emotions, and overall health during pregnancy. Conditions for disturbed sleep may arise because of hormonal shifts, nausea, frequent urination, backache, cramps in the legs, heartburn, movements of the fetus, or feelings of anxiety. Good sleep habits involve making sure of a regular time for going to bed and for waking up, creating the right sleeping conditions, limiting the use of caffeine, particularly at late hours, avoiding too heavy meals shortly before going to bed, and minimizing active work just before going to bed. The right sleeping position and the use of special pillows would help promote sleep at every stage of pregnancy. If insomnia is constant, or if one feels rather sleepy during the day, it is recommended to see a doctor. [32]

3.8.4 Social and Emotional Support

Sufficient emotional and social support helps women cope with stress and strains of pregnancy. Family members, friends, partners, social groups, and caregivers can provide emotional comfort and useful help. Family and friends offer support through assistance with domestic chores, encouragement of healthy diet and exercise, accompaniment on visits to healthcare facilities, and active listening. Caregivers are supposed to communicate in a polite and encouraging way and to stimulate girls' discussions about their emotional problems during the appointments. In case any serious psychological problem is detected on time, a referral to a relevant specialist is very beneficial for obtaining proper treatment and support. [33]

3.9 Lifestyle Modification for Safe Motherhood

As mentioned in (Table 3.3) the role of healthy habits during pregnancy is critical in promoting maternal wellness, enabling normal fetal growth and development, and decreasing the chances of having problems related to pregnancy. There are many changes in physiology and behavior during pregnancy, which is why this moment is very important to determine the healthy patterns related to nutrition, physical activities, weight control, sleeping habits, avoiding bad habits, keeping hygiene, and using medications. Lifestyle advice should be personalized and depend on pregnancy time, body mass index before pregnancy, nutrition state, level of activity, cultural habits, and the presence of some diseases or complications associated with the pregnancy. It is

necessary to receive constant counseling and monitor pregnant women for helping them to make better choices and maintain healthy behavior during pregnancy. [34]

3.9.1 Healthy Weight Gain

Correct amount of weight gain during pregnancy serves as an essential measure of nutritional health of the woman and pregnancy status. The amount of advisable weight gain is closely related with the weight of the mother before pregnancy, with ranges applied to women, depending whether she was seen as underweight, normal, overweight or obese before pregnancy, thus ranging weight gain recommendations. Less than recommended weight gain can lead to risk of slow fetal growth and low birth weight. Excessive weight gain can lead to risks of developing gestational diabetes, complications related to high blood pressure, need for cesarean delivery or retaining unnecessary weight after pregnancy. The weight of the expectant mother must be carefully monitored during pregnancy and her diet and also physical activity adjusted accordingly. [35]

3.9.2 Tobacco and Alcohol Avoidance

To ensure both maternal and fetal health, it is important to avoid tobacco, nicotine and alcohol. Smokers put their fetus at risk of harmful substances and chemicals, including nicotine and carbon monoxide. These chemicals will affect the availability of oxygen for the fetus and its growth. Maternal smoking or second-hand smoking has been linked to multiple outcomes such as low birth weight, preterm birth, placental abnormalities, fetal growth restriction and stillbirth. The pregnant women should be encouraged to stop smoking at the earliest possible time and avoid tobacco smoke exposure at all costs, as well-assisted behavioral advice and strategies will help if necessary. Alcohol consumption should be completely avoided as well, as prenatal exposure to this substance may impede fetal growth and development thus leading to long-term negative effects.

3.9.3 Caffeine and Other Dietary Considerations

Caffeine passes through the placenta, and the fetus’s ability to break down caffeine is limited. High caffeine intake in mothers has been linked to negative consequences in the pregnancy, such as higher chance of pregnancy loss and lower birth weight. Therefore, it should be kept in mind that pregnant women should limit the caffeine intake according to commonly known guides and recommendations, keeping in mind the total caffeine intake from beverages and diets (i.e. coffee, tea, cola, energy drinks, chocolate food and medication). The diet must consist mostly of vegetables, fruits, cereals, legumes, dairy products, eggs, lean meat, nuts, and seeds. Foods rich in sugar, sodium, saturated fats, and processed ingredients must be used minimally.

3.9.4 Food Safety and Hygiene

Pregnancy causes changes in the immune system that make pregnant women more susceptible to food-borne infections. Bacteria like Listeria monocytogenes, Salmonella, Toxoplasma gondii, and Campylobacter can result in increasingly severe complications for mothers and their babies. Pregnant women are advised to refrain from eating raw or undercooked fish, poultry, meat, or eggs. Furthermore, unpasteurized milk and dairy products must also be avoided pregnant women. All fruits and vegetables should be thoroughly washed. It is important to note that hands need to be washed frequently and that proper food preparation requires the use of sanitary methods and practices, including clean refrigerators and methods of cooking. Even though fish is a good source of protein and omega 3 fats, consumption of fish should be limited due to mercury contamination.

3.9.5 Healthy Sleep and Daily Routine

Sufficient sleep as well as adequate rest is essential for health, safety, and effectiveness of operations while being pregnant. Disorders sleep may arise due to physiological changes, such as hormonal processes, nausea, frequent urination, back pain, cramping of legs, heartburn, foetal movements, or anxiety. A pregnant woman should keep a regular bed-time and wake-up schedule and make her sleeping environment calm and warm. To sleep better, a pregnant woman must limit the intake of caffeine, eliminate heavy meals before going to bed, and avoid active leisure before sleeping. Physical activities during the day and relaxation exercises may help sleep soundly. Regular meals, enough fluids, sufficient physical activities, preventing overload, and sufficient time for resting should be considered when planning a daily schedule. In case of severe disturbances of sleeping, the advice of a doctor should be sought.

3.9.6 Medication and Supplement Safety

Special care should be given to the use of drugs or supplements during pregnancy since certain medication can cross the placenta and thus affect the developing fetus. Women who plan on taking medicines, either prescription or over-the-counter, supplements, or herbal products during pregnancy must seek medical advice. The health care provider should review the women's past medical history, weighing the beneficial effects to the mother and possible risk of the baby. No medication should be stopped without the doctor's approval since untreated conditions of the mother may pose a risk to pregnancy. Certain supplements may be recommended depending on the nutritional needs of the pregnant woman. Women should not self-medicate and refrain from excessively taking supplements or using unverified herbal products.

Overall, making lifestyle changes for a risk-free pregnancy entails a multifaced and personalized strategy that includes proper weight gain during pregnancy, healthy eating habits, physical activity, abstaining from cigarette and alcohol use, proper caffeine consumption, food

safety practices, adequate sleep and rest, as well as proper medication and supplement consumption. Ongoing education during antenatal visits can help pregnant women detect possible dangers, learn about healthy habits, and make wise choices. These actions together with regular antenatal healthcare and good management of pregnancy complications will ensure a better health of mothers and a safer pregnancy experience.

Table 3.3: Healthy Lifestyle Practices for a Safe Pregnancy

Lifestyle Factor	Recommended Practice	Potential Benefit
Healthy weight gain	Monitor weight regularly according to pre-pregnancy BMI	Supports appropriate fetal growth and reduces pregnancy-related complications
Balanced diet	Consume a varied, nutrient-dense diet including whole grains, fruits, vegetables, legumes, dairy, eggs, nuts and lean protein	Supports maternal health and fetal development
Physical activity	Engage in regular, moderate physical activity when medically appropriate	Supports fitness, healthy weight gain and psychological well-being
Tobacco and alcohol	Avoid tobacco, nicotine, second-hand smoke and alcohol	Reduces risks to fetal growth and development
Caffeine	Limit total caffeine intake from coffee, tea, cola, energy drinks, chocolate and medicines	Helps minimize potential adverse pregnancy effects
Food safety	Avoid raw/undercooked foods and unpasteurized products; wash produce and maintain proper food hygiene	Reduces risk of foodborne infections
Sleep and rest	Maintain regular sleep and adequate periods of rest	Supports energy, emotional well-being and overall health
Hydration	Maintain adequate fluid intake throughout pregnancy	Supports blood volume, amniotic fluid and normal physiological functions
Medication and supplements	Use prescription, OTC medicines, supplements and herbal products only after appropriate medical advice	Helps minimize potential fetal and maternal risks
Antenatal care	Attend regular antenatal visits and receive individualized counselling	Allows monitoring, early identification of problems and timely intervention

3.10 Role of Healthcare Professionals

Providers of health care and medicine play an important role in promoting wellness, nutrition, fitness and mental well-being, and providing safe living recommendations to pregnant women. All relevant parties including midwives, obstetricians, physicians, nurses, pharmacists, dietitians and others can provide customized nutrition education and consider previous lifestyle issues that have occurred. This coordinated approach ensures that the needs of the pregnant woman as well as those of her unborn baby will be met throughout the entire pregnancy.

3.10.1 Nutritional Counselling

Nutritional counselling should start early in pregnancy and continue during the antenatal period. Evaluation of diet, pre-pregnancy BMI, weight gain during pregnancy, preferences, nutrition-related risks, and diseases should be done by a healthcare professional. Pregnant women should be informed about what balanced meals are, how much food, energy and protein should be consumed and the importance of vitamins and minerals, hydration, food safety and supplementation. It should be noted that counselling should be practical and personalized regarding food culture, food availability, sociocultural and economic factors and personal tastes. Women at risk for nutritional deficiencies and stress-related pregnancy complications may require more extensive evaluation and treatment by a dietitian or other specialist.

3.10.2 Exercise and Lifestyle Counselling

Based on their health condition, women should receive recommendations from qualified healthcare personnel regarding correct exercise and healthy lifestyle behavior. Women with no contraindications should be advised moderate exercise, which is good for physical condition, healthy weight gain during pregnancy, and general health. Advice must include appropriate types of physical activity with proper intensity, hydration, warning signs and contraindications for physical activity. Healthcare specialists should also give recommendations about quitting smoking and drinking alcohol, reduced caffeine intake, sufficient sleep, stress relief, food hygiene, and proper medications. Individualized recommendations have a great importance for those women that are suffering from medical or pregnancy complications.

3.10.3 Monitoring Maternal and Fetal Health

Consistent prenatal supervision allows clinicians to monitor the health of both mother and fetus and eliminate any potential complications. As far as mother's health is concerned, healthcare professionals must monitor such vital indicators as weight gain, blood pressure, levels of nutrients, as well as other laboratory parameters including haemoglobin level. The assessing of fetal health and development will include following parameters: growth of fetus, heart rate, and indicators of development according to other clinical measures. In addition to that, healthcare personnel must monitor adherence to medicinal prescriptions aimed at promoting a healthy

lifestyle. Any abnormal results must be investigated and managed. Communication between members of the healthcare team is of great importance.

3.11 References

1. Kartchner LC, Dunn A, Taylor KH, Ali MM, Manning NA, Dajani NK, Magann EF. Lifestyle modifications prior to pregnancy and their impact on maternal and perinatal outcomes: a review. Journal of Clinical Medicine. 2025 Sep 18;14(18):6582.
2. Habibi F, Shahhosseini Z, Jahanfar S, Fendereski A, Ahmadi M, Hamzehgardeshi Z. Promoting a healthy lifestyle during pregnancy in Iran: a best practice implementation project. JBI Evidence Implementation. 2026 Jun 22:10-97.
3. Procter SB, Campbell CG. Position of the Academy of Nutrition and Dietetics: nutrition and lifestyle for a healthy pregnancy outcome. Journal of the Academy of Nutrition and Dietetics. 2014 Jul 1;114(7):1099-103.
4. Farpour-Lambert NJ, Ells LJ, Martinez de Tejada B, Scott C. Obesity and weight gain in pregnancy and postpartum: an evidence review of lifestyle interventions to inform maternal and child health policies. Frontiers in endocrinology. 2018 Sep 26;9:546.
5. Ainscough, K.M., O'Brien, E.C., Lindsay, K.L., Kennelly, M.A., O'Sullivan, E.J., O'Brien, O.A., McCarthy, M., De Vito, G. and McAuliffe, F.M., 2020. Nutrition, behavior change and physical activity outcomes from the PEARS RCT—An mHealth-supported, lifestyle intervention among pregnant women with overweight and obesity. Frontiers in endocrinology, 10, p.938.
6. Albracht-Schulte KD, García-González Á, Wilson S, Robert-McComb JJ. Nutritional guidelines and energy needs during pregnancy and lactation for active women. InThe active female: health issues throughout the lifespan 2023 Feb 26 (pp. 363-378). Cham: Springer International Publishing.
7. Khammarnia M, Ansari-Moghaddam A, Kakhki FG, Clark CC, Barahouei FB. Maternal macronutrient and energy intake during pregnancy: a systematic review and meta-analysis. BMC Public Health. 2024 Feb 15;24(1):478.
8. Lezhneva JO, Selezneva IS. NUTRIENT REQUIREMENTS DURING WOMEN PREGNANCY AND LACTATION. The School on Biotechnology for Students, Ph. D. students and Young scientists.—Yekaterinburg, 2025. 2025:89-92.
9. Fatima N, Yaqoob S, Rana L, Imtiaz A, Iqbal MJ, Bashir Z, Shaukat A, Naveed H, Sultan W, Afzal M, Kashif Z. Micro-nutrient sufficiency in mothers and babies: management of deficiencies while avoiding overload during pregnancy. Frontiers in Nutrition. 2025 Apr 1;12:1476672.
10. Song Y, Zhang F, Lin G, Wang X, He L, Li Y, Zhai Y, Zhang N, Ma G. A study of the fluid intake, hydration status, and health effects among pregnant women in their second trimester in China: A cross-sectional study. Nutrients. 2023 Apr 2;15(7):1739.

11. Song Y, Zhang F, Wang X, Lin G, He L, Lin Z, Zhang N, Ma G. A study of fluid intake, hydration status, and body composition of pregnant women in their third trimester, and relationships with their infant's birth weight in China: a prospective cohort study. Nutrients. 2024 Mar 27;16(7):972.
12. Nayak R, Agrawal N, Chandravanshi L, Rathia S, Kurre P. Ayurvedic Concept of Garbhini Paricharya: A Critical Review. Pharmacognosy Reviews. 2025;19(37):30-8.
13. McKay FH, Spiteri S, Zinga J, Sulemani K, Jacobs SE, Ranjan N, Ralph L, Raeburn E, Threlfall S, Bergmeier ML, van der Pligt P. Systematic review of interventions addressing food insecurity in pregnant women and new mothers. Current nutrition reports. 2022 Sep;11(3):486-99.
14. Asim M, Ahmed ZH, Nichols AR, Rickman R, Neiterman E, Mahmood A, Widen EM. What stops us from eating: a qualitative investigation of dietary barriers during pregnancy in Punjab, Pakistan. Public health nutrition. 2022 Mar;25(3):760-9.
15. Maggiulli O, Rufo F, Johns SE, Wells JC. Food taboos during pregnancy: meta-analysis on cross cultural differences suggests specific, diet-related pressures on childbirth among agriculturalists. PeerJ. 2022 Jul 11;10:e13633.
16. Zych-Krekora K, Sylwestrzak O, Krekora M. The critical role of iron in pregnancy, puerperium, and fetal development. Journal of Clinical Medicine. 2025 May 16;14(10):3482.
17. Gomes F, Ashorn P, Askari S, Belizan JM, Boy E, Cormick G, Dickin KL, Driller-Colangelo AR, Fawzi W, Hofmeyr GJ, Humphrey J. Calcium supplementation for the prevention of hypertensive disorders of pregnancy: current evidence and programmatic considerations. Annals of the New York Academy of Sciences. 2022 Apr;1510(1):52-67.
18. Prasad HA, Sylvester SK. Nutritional Deficiencies and Requirements during Pregnancy. Asian Journal of Applied Science and Technology (AJAST). 2025 Jan;9(1):129-44.
19. Thompson B, Amoroso L. Combating micronutrient deficiencies: food-based approaches. 2010 Dec 22.
20. Olloqui-Mundet MJ, Cavia MD, Alonso-Torre SR, Carrillo C. Dietary habits and nutritional knowledge of pregnant women: the importance of nutrition education. Foods. 2024 Oct 8;13(19):3189.
21. Alamri SH, Abdeen GN. Maternal nutritional status and pregnancy outcomes post-bariatric surgery. Obesity Surgery. 2022 Apr;32(4):1325-40.
22. Cilar Budler L, Budler M. Physical activity during pregnancy: a systematic review for the assessment of current evidence with future recommendations. BMC Sports Science, Medicine and Rehabilitation. 2022 Jul 16;14(1):133.
23. Paulsen CP, Bandak E, Edemann-Callesen H, Juhl CB, Händel MN. The effects of exercise during pregnancy on gestational diabetes mellitus, preeclampsia, and spontaneous abortion

among healthy women—a systematic review and meta-analysis. International Journal of Environmental Research and Public Health. 2023 Jun 6;20(12):6069.
24. Yang X, Li H, Zhao Q, Han R, Xiang Z, Gao L. Clinical practice guidelines that address physical activity and exercise during pregnancy: a systematic review. Journal of midwifery & women's health. 2022 Jan;67(1):53-68.
25. Szumilewicz A, Santos-Rocha R. Exercise selection and adaptations during pregnancy. InExercise and Physical Activity During Pregnancy and Postpartum: Evidence-Based Guidelines 2022 Oct 27 (pp. 275-361).
26. Hayman M, Brown WJ, Brinson A, Budzynski-Seymour E, Bruce T, Evenson KR. Public health guidelines for physical activity during pregnancy from around the world: a scoping review. British Journal of Sports Medicine. 2023 Jul;57(14):940-7.
27. Kuder L, Dinevski D, Dinevski IV, Takač I, Mujezinović F, Gašparović VE. Benefits of yoga in pregnancy: a randomised controlled clinical trial. Journal of Perinatal Medicine. 2025 Mar 1;53(3):350-7.
28. Villar-Alises O, Martinez-Miranda P, Martinez-Calderon J. Prenatal yoga-based interventions may improve mental health during pregnancy: an overview of systematic reviews with meta-analysis. International Journal of Environmental Research and Public Health. 2023 Jan 14;20(2):1556.
29. Saini T, Panwar A. Benefits of yoga during pregnancy. Wide Vision. 2023 Jun 22;151.
30. Penner F, Rutherford HJ. Emotion regulation during pregnancy: a call to action for increased research, screening, and intervention. Archives of women's mental health. 2022 Apr;25(2):527-31.
31. Jalal SM, Alsebeiy SH, Alshealah NM. Stress, anxiety, and depression during pregnancy: A survey among antenatal women attending primary health centers. InHealthcare 2024 Nov 7 (Vol. 12, No. 22, p. 2227). MDPI.
32. Meers JM, Nowakowski S. Sleep during pregnancy. Current psychiatry reports. 2022 Aug;24(8):353-7.
33. Al-Mutawtah M, Campbell E, Kubis HP, Erjavec M. Women's experiences of social support during pregnancy: a qualitative systematic review. BMC pregnancy and childbirth. 2023 Nov 10;23(1):782.
34. Kartchner LC, Dunn A, Taylor KH, Ali MM, Manning NA, Dajani NK, Magann EF. Lifestyle modifications prior to pregnancy and their impact on maternal and perinatal outcomes: a review. Journal of Clinical Medicine. 2025 Sep 18;14(18):6582.
35. Makama M, Skouteris H, Moran LJ, Hill B, Redman LM, Lim S. Co-designing a community lifestyle intervention program to reduce postpartum weight retention. Health Expectations. 2024 Feb;27(1):e13905.