WISELEAF SCIENTIFIC VENTURES



Chapter 4: Antenatal Care, Investigations and Preventive Healthcare during Pregnancy

Prof. Jagruti R. Patil

Assistant Professor, Department of Pharmaceutics, R V P M Institute of Pharmacy, Dwarka, Nashik, India.

Mail id – rxutkarshmandage@gmail.com

Abstract

Antenatal care is one of the most important aspects of maternal health care. This service ensures that mothers and their babies remain healthy throughout their pregnancy. Antenatal visits allow doctors to monitor mothers' physiological changes, assess the state of the fetus, detect high-risk pregnancies, and identify complications before they pose any serious threat. In this chapter, the major principles and schedule of antenatal care are provided, taking into account the importance of regular examinations of both mothers and babies. It describes the most significant laboratory investigations, such as blood tests, blood group and Rh factor identification, glycosuria, and others. It further clarifies the importance of ultrasound imaging, fetal heart rate measurement, and various other techniques employed in evaluating the development of the fetus. The chapter discusses the importance of educating patients about different frameworks related to pregnancy, like nutrition, treatment regimen, and dangers during pregnancy, childbirth, breastfeeding, as well as mental health and family assistance.

Keywords

Antenatal Care; Maternal Health; Fetal Monitoring; Preventive Healthcare; Pregnancy Vaccination

4.1 Introduction

Pregnancy can be characterized as a complex physiological condition which results in the occurrence of important anatomical and hormonal modifications as well as metabolic and cardiovascular changes in the mother. These modifications are important for the development of the placenta, development of the fetus, and preparation for childbirth and breastfeeding. Although pregnancy is regarded a normal physiological process in the majority of women,

many complications can arise at any point in time, and such issues may impact the health condition of both women and babies. Conditions which include anemia, gestational diabetes mellitus, pregnancy-related high blood pressure, infections, issues with the growth of the fetus, and others may appear undetected in the first stage of pregnancy. Thus, systematic antenatal care is vital for ensuring the good condition of the mother, monitoring the fetus, and identifying possible complications as soon as possible. [1]

Antenatal care entails health services offered to women during pregnancy through a prearranged series of encounters with qualified health personnel. The specific aims of antenatal care are to protect the health of the mother and the fetus, analyze the course of pregnancy, recognize risk factors, avert possible complications, and ensure appropriate and timely treatment when needed. When undergoing antenatal visits, women may receive information regarding their medical and obstetric health history, be examined medically, their vital parameters monitored, undergo an assessment of their health and nutritional condition, and the fetal development process. Antenatal visits may differ in terms of frequency and content depending on the particular stage of pregnancy, health condition of the pregnant woman, as well as pregnancy risk. [2]

It is essential to schedule regular antenatal appointments since they help in the process of continuous monitoring and not depending on just a one-time pregnancy evaluation. Maternal weight, blood pressure, symptoms, and other related clinical findings are tracked over time, allowing health specialists to detect any deviation from the expected changes during pregnancy. Fetal assessment is carried out through measuring fetal growth or monitoring fetal heartbeat and through imaging and several other techniques. As a result, regular appointments provide an opportunity to communicate tips on healthy diet, physical exercises, and compliance with medications, taking supplements, vaccination, and other preventive measures. Laboratory tests play an important role in the antenatal assessment process. Common tests conducted during pregnancy include CBC, Hb tests, typing of blood groups and Rh factor, urine tests, blood sugar tests, and tests for specific infections. Along with these tests, other tests like thyroid, kidney, liver, or other tests may also be performed when necessary. Early detection of abnormalities is important in order to act appropriately and achieve good results concerning both the pregnancy and the mother. [3]

Ultrasonography and fetal monitoring provide data on the condition of the fetus, gestational age, anatomy, and growth of the fetus as well as location of the placenta. Different types of ultrasound examinations may be carried out during different periods of pregnancy depending on clinical indications. For women with high-risk pregnancies, extra fetal tests such as Doppler tests, cardiotocography, non-stress tests, and biophysics can be required. Preventive healthcare is a crucial part of antenatal care, as carrying out proper nutrition supplements such as iron, calcium, and folic acid helps in the avoidance of nutrition deficiency and its consequences.

Maternal vaccination protects the mother and sometimes also gives the newborn passive immunity. Infection prevention, personal hygiene, food safety, proper physical exercise and rest, and the avoidance of tobacco, alcohol, or any other harmful substances form part of good practices for healthy pregnancy. [4]

Patient education and counsellig should form part of every antenatal visit as pregnant women and their families must get accurate and timely information about nutrition, medications and supplements, tests, vaccinations, physical activity, warning signs, birth planning, and where to go for medical help in case of emergencies. Counselng should be tailored according to the needs of women at risk and taking into consideration their health condition and cultural differences. Antenatal care gives us the chance to discover the women who need referral to specialized services and consultations. Pregnancies complicated by pre-existing health issues, previous negative results in pregnancy, twin pregnancies, abnormal fetal conditions, etc. may need to receive appropriate care and timely referral. Multidisciplinary cooperation of obstetricians, nurses, midwives, pharmacists, nutrition specialists, etc. can help in delivering complete and patient-oriented services.

Antenatal care covers the full cycle of preventive and diagnostic activities including preventive and supportive aspects aimed at healthy pregnancy. A comprehensive approach including scheduled antenatal visits, evaluation of the mother/fetus, the necessary laboratory tests, vaccination, preventive care measures, and counseling allows one to detect complications early and improve both maternal and fetal health. This paper highlights some components of antenatal care, types of medical tests and procedures, fetal monitoring, vaccination and preventive measures, etc.

4.2 Antenatal Care (ANC)

Antenatal care (ANC) is defined as the systematic healthcare provided to women in the time leading up to delivery in order to facilitate maternal and fetal well-being and support a healthy pregnancy. It involves following the woman during her pregnancy through routine assessments of her well-being, assessment of risk factors, preventive measures, required tests, treatment of diseases, and health education. ANC is a means of caring for pregnant women which gives health professionals the opportunity to ensure the normal course of pregnancy and detect any problems at its early stages.

4.2.1 Objectives and Principles

The aims of antenatal care (ANC) are safe maintenance of the mother and the fetus, following the mother's progress throughout her pregnancy, recognizing problems as soon as they appear, and providing them advice or treatment in a timely manner. The recommendations made to the mother deal with the evaluation of her weight, blood pressure, nutrition, and any clinical symptoms, as well as the safety of her unborn baby based on the gestational age. In addition,

ANC is a chance for undergoing different tests. It also offers a number of prevention methods such as nutritional supplementation and vaccinations. Finally, ANC provides the mother with information about medicine, risks of pregnancy, and checkup scheduling. [5]

4.2.2 ANC Schedule

ANC should start in the first trimester or immediately after pregnancy confirmation. The first visit will typically be to confirm and date the pregnancy, collect full medical and obstetric histories; conduct physical examination; evaluate the nutritional state and risks; carry out baseline investigations and commence essential preventive care. According to World Health Organization, it is recommended that there be at least eight ANC visits throughout the pregnancy to allow for good continuity of care. Visits thereafter will be to monitor the health of the fetus and mother, check the results of investigations, determine any new abnormal symptoms, and provide preventive interventions. Women with high-risk pregnancies, abnormal findings, and pre-existing medical disorders will require more frequent visits. [6]

4.2.3 Trimester-wise Care

As mentioned in (Table 1) Antenatal care is tailored to meet the physiological changes that occur during each trimester. The first trimester is centered on determining whether a woman is pregnant, calculating how many weeks pregnant she is, checking the health and history of the mother, carrying out preliminary tests and starting taking folic acid and preventive drugs. There is also counseling on diet, treatments, lifestyle, and avoidance of risks. In the second trimester, the care given to the pregnant woman involves the continual monitoring of both maternal and fetal health, carried out in accordance with the required assessments. The screening method that is used to determine conditions like anemia, gestational diabetes, diseases, and fetal abnormalities takes into consideration the individual's risk factors and the recommendations. An ultrasound might be conducted in order to monitor the fetus's anatomical and developmental stages. The care provided during the final trimester involves the monitoring of the fetal growth and assessing the well-being of the mother. Routine assessments will be carried out during this stage to evaluate the mother's condition and the baby's development too. In addition, the doctor provides special information that the mother will need about labor and delivery, labor signs, actions to take during emergencies, breastfeeding, and newborn care. [7]

Table 1: Trimester-wise Care during Pregnancy

Trimester	Major Areas of Care	Key Components
First Trimester	Initial assessment and preventive care	Confirmation and dating of pregnancy; maternal history and health assessment; preliminary investigations; folic acid and preventive medications;

		counselling regarding diet, treatment, lifestyle and avoidance of risks
Second Trimester	Maternal and fetal monitoring	Continuous monitoring of maternal and fetal health; screening for anemia, gestational diabetes, infections and fetal abnormalities according to risk factors; ultrasound assessment of fetal anatomy and development when indicated
Third Trimester	Fetal growth and maternal well-being	Monitoring fetal growth and maternal condition; routine maternal and fetal assessments; counselling regarding labour and delivery, signs of labour, emergency measures, breastfeeding and newborn care

4.2.4 High-Risk Pregnancy Identification

A pregnancy is said to be high-risk if the mother or the fetus is prone to complications or health issues during the pregnancy, childbirth, or postpartum. Therefore, the risk factors should be identified during the first antenatal care appointment and continuously during the entire course of the pregnancy since new problems can appear at any time. [8] Risk factors include the mother's age, either extremely young or old, prior poor pregnancy outcomes, multiple pregnancy, existing diseases such as diabetes and hypertension, as well as complications during pregnancy, such as diabetes, hypertension, noteworthy anemia, bleeding, etc. Expectant mothers who are classified as having a high-risk pregnancy need personalized evaluation and treatment that is determined by the individual risk factors. They may require extra prenatal checkups, additional lab tests, more advanced ultrasound, obstetric monitoring, therapy changes, meeting with specialists, or referral to the correct institution. Appropriate treatment and early identification of risk factors allows clinicians to apply needed measures. [9]

4.3 Importance of Regular Check-ups

Regularly scheduled visits to the doctor are an essential component of pregnancy-related medical care. Unlike one-time reviews, repeated inquiries make it possible for physicians to observe changes or issues over time, compare those changes against standards of normal pregnancy, and detect problems early in pregnancy. During each visit, the doctor is able to get information about the mother's health condition, the state of the fetus and the results of medical tests conducted on the mother. Regular visits also help keep one healthy, solve many problems, and provide advice and treatment whenever necessary.

4.3.1 Maternal Health Monitoring

Regular antenatal checkups can evaluate the mother's physical, nutritional, and psychological condition throughout her pregnancy. [10] Changes in the mother's weight, blood pressure, pulse rate, and other health indicators are monitored at prescribed intervals. Doctors observe for

symptoms of severe nausea, headaches, dizziness, abdominal discomfort, vaginal bleeding or swelling, temperature increase, and any urinary disorders. One can also check the mother's nutritional health and adherence to dietary supplements like iron, folic acid, and calcium. Additionally, routine checkups can also reveal specific diseases, such as anemia, gestational diabetes, hypertension, thyroid dysfunction, or any infection. Women with pre-existing medical problems can get special care as necessary. [11]

4.3.2 Fetal Well-being Monitoring

Fetal well-being should be continuously reviewed in order to ascertain fetal development. When receiving prenatal care, fetal heart activity will be recorded and is evaluated based on the age of the while uterine growth and other factors can indicate fetal well-being. Use of ultrasound will be performed at certain points of gestation to evaluate gestational age and fetal anatomy as well as obtain details of placental situation, amniotic fluid, etc. [12] Specific cases may also involve additional tests such as Doppler tests, non-stress tests, CTG, etc. Monitoring fetal movements during the last trimester can yield important information about fetal health and can lead to discovery of any problems. [13]

4.3.3 Early Detection of Complications

One of the key advantages of antenatal follow-up is its ability to aid in the early identification of severe complications. Through various forms of screening and clinical assessment together with timely investigations we can possibly detect such medical conditions as anemia, gestational diabetes, hypertensive disorders, infections and anomalies in fetal growth and other maternal and fetal diseases before they become severe. [14] Early identification of conditions such as anemia allows for timely treatment, monitoring patient condition closely, and if required changing medications and consulting specialists. Another important feature of regular antenatal visits is the possibility of educating women on the signs that need urgent medical attention. Women should learn to identify such urgent symptoms as severe headache, visual disturbances, vaginal bleeding, loss of amniotic fluid, abdominal pain, fever, difficulty breathing, convulsions, or decreased fetal activity. [15]

4.4 Laboratory Investigations

As mentioned in (Table 2) Investigations done in a laboratory are a crucial part of prenatal care and play a role in determining the health status of the mother, detecting pregnancy-related conditions, and identifying disorders which may affect the health of the mother or baby. The tests are typically done during the first prenatal visit in accordance with the same local guidelines and different stages of pregnancy depending on gestational age, clinical findings and risk factors. Some of the tests commonly has done include blood tests, blood grouping and Rh typing, urinalysis, glucose test, screening for infectious diseases, and other tests if needed.

Early detection of health conditions ensures early treatment, hence increasing chances of good pregnancy outcomes. [16]

4.4.1 Hematological Investigations

The primary aim of performing hematological tests is to examine the blood of the mother and detect conditions like anemia and various other hematological disorders. A complete blood count (CBC), which is also referred to as a blood test, is usually included in the investigation, which consists of hemoglobin levels, hematocrit, red blood cell indices, white blood cell count and platelet count. The determination of the hemoglobin level is one of the most important tests because iron deficiency anemia is often found in pregnant women and can lead to other issues such as exhaustion and low birth rate or premature birth. Additionally, the investigation of red blood cell indices can help understand the type of anemia while evaluating. Platelet counts can identify the presence of thrombocytopenia, which might serve as an important indicator for assessing some other pregnancy-related complications. Depending on clinical findings, some other blood tests like peripheral blood smear, serum ferritin and other hematological tests might be performed. [17]

4.4.2 Blood Group and Rh Typing

A key aspect of the initial assessment during antenatal care is determining the mother's blood group and Rh status. The need to identify an Rh-negative mother is underscored by the fact that the mother's sensitization to Rh-positive cells can lead to hemolytic disease of a future or concurrent pregnancy. If necessary, the further analysis of red blood cell antibodies using an antibody screen may be performed to find women who have undergone alloimmunization. With the help of this information, a strategy for its monitoring and preventive treatment can be formulated. Moreover, knowing the blood group is also helpful in forming a strategy for a blood transfusion in case the latter has to happen. [18]

4.4.3 Urine and Glucose Testing

Urine testing is a common procedure performed during pregnancy for detection of the possibility of abnormalities in urine and diagnoses certain types of health conditions. The normal urinalysis routine provides some essential information regarding the presence of protein, glucose, blood, leukocyte, and nitrites. Detection of proteinuria, especially in cases when the patient has elevated blood pressure may indicate the need for further tests on pregnancy-related hypertension. Urinary symptoms may signal the need for urine culture test for establishment of diagnosis regarding urinary tract infection. Glucose level analysis is done to determine hyperglycemia and to diagnose gestational diabetes mellitus (GDM). Usually done in the trimester prescribed, screening can be performed earlier in women who are identified as being at risk. The screening may involve glucose challenge tests or a glucose tolerance test depending on the protocols being put in place. Detection of GDM early enough

and putting into place measures like dieting, engaging in physical activities, monitoring blood glucose levels, and using medication where needed has been shown to minimize risks to the mother and the baby. [19]

4.4.4 Infectious Disease Screening

It's important to screen specific infectious diseases during pregnancy because maternal infections can have negative effects on the mother, fetus, or newborn, and can sometimes be transmitted either during pregnancy or delivery. Generally, antenatal screening includes tests for HIV, hepatitis B infection, and syphilis, but the exact screening panel varies depending on national recommendations and each individual's risk factors. Early detection makes it possible to offer effective treatment or preventive measures, thus minimizing the chances of transmission to the newborn. Other infections that can potentially be screened include hepatitis C, TB, rubella immunity, or some other infections that are a priority in a certain geographic region depending on clinical background, epidemiological risk level, vaccination status, etc. [20]

4.4.5 Other Investigations

Further laboratory studies might be advised based on information taken from the woman's past, her symptoms, examination results and risk indicators for pregnancy. If existence of the disease of the thyroid is suspected, testing for its functioning will be needed. Women may need tests of liver and kidney functioning if they have high blood pressure, previously existing kidney or liver disease, or indications of developing complications related to pregnancy. Testing for coagulation may be needed in some cases when there is a suspicion of bleeding disorders as well as complications during pregnancy. Different investigations including testing for certain infections, nutrition deficiency or autoimmune diseases may also be necessary. At the same time decisions about the necessity of conducting additional investigations must be made based on individual approach instead of standardizing the testing for every woman. Overall, laboratory testing in antenatal conditions allows the doctors to gather essential information about the health of the mother and xreate a list of possible diseases that might require specific medical interventions in future.

Table 2: Common Laboratory Investigations during Pregnancy and Their Purpose

Laboratory Investigation	Primary Purpose	Major Conditions/Findings Identified
Complete blood count (CBC)	Assessment of overall hematological status	Anemia, thrombocytopenia and other hematological abnormalities
Hemoglobin estimation	Assessment of maternal hemoglobin level	Detection and monitoring of anemia

Blood group and Rh typing	Determination of ABO and Rh status	Identification of Rh incompatibility and risk of alloimmunization
Urinalysis	Assessment of urinary abnormalities	Proteinuria, glycosuria, hematuria and urinary infection-related findings
Blood glucose testing	Assessment of maternal glucose status	Detection of hyperglycemia and gestational diabetes mellitus
HIV screening	Screening for maternal HIV infection	Identification of HIV infection and need for appropriate preventive measures
Hepatitis B screening	Screening for hepatitis B virus infection	Identification of maternal HBV infection
Syphilis screening	Screening for maternal syphilis	Detection of syphilis requiring treatment
Thyroid function tests	Assessment of thyroid function when clinically indicated	Thyroid dysfunction
Liver function tests	Evaluation of hepatic function when indicated	Liver dysfunction and pregnancy-related hepatic abnormalities
Renal function tests	Assessment of kidney function when indicated	Renal impairment and related abnormalities
Peripheral blood smear	Further evaluation of abnormal blood findings when indicated	Morphological abnormalities and selected causes of anemia
Serum ferritin	Assessment of iron stores when indicated	Iron deficiency and depleted iron stores

4.5 Ultrasonography and Fetal Monitoring

Ultrasound and fetal monitoring are very useful aspects of prenatal care that help assess the fetal growth and condition. As shown in (Fig 1) Ultrasound provides images of the fetus and adjacent structures without using radiation for the mother or the child. The type of ultrasound used and the time when this ultrasound is performed depend on the term of pregnancy, characteristics of the pregnancy, clinical indications, and national recommendations. Fetal monitoring implies analysis of the fetal heart rate, movements, development, quantity of amniotic fluid, and placental situation and even blood circulation in the uteroplacental and fetoplacental systems on request. All these evaluations allow finding out the pregnancies that should be watched after more carefully and treated at the hospital.

4.5.1 Ultrasound Scans

Ultrasound is a commonly practiced examination in pregnancy which helps in determining fetal viability, gestational age, and growth, as well as the anatomy and location of the fetal placenta, etc. An ultrasound test is usually performed at an early stage of pregnancy to ensure intrauterine pregnancy, predict viability, determine gestational age and identify multiple pregnancies. An ultrasound examination during the first trimester may also help in screening for certain fetal chromosomal abnormalities if used with other maternal assessments. The mid-trimester ultrasound, usually performed at established periods of gestation in accordance with local guidelines, gives a thorough assessment of the fetal anatomy and detects many structural defects. Further ultrasound tests can be done when necessary, in order to evaluate fetal growth and amniotic fluid amount and ensure fetal position and presentation, etc. Thus, ultrasound has proven to be very helpful in detecting some cases like suspected fetal growth restrictions, multiple pregnancies, abnormal fundal heights, etc. [21]

4.5.2 Fetal Growth and Heart Rate Monitoring

Observing the baby's growth is crucial in pregnancy care, as abnormal growth may lead to placental problems, maternal diseases, nutritional issues, or other problems associated with pregnancy. The clinical evaluation can include the evaluation of fundal height and growth of the fetus. In case of suspected problems with development, ultrasound can be used to estimate the size of the fetus and evaluate development over time. Routine evaluations can be completed at visits made if growth problems are suspected, or risk factors of fetal growth are evident. The experience of fetal heart performance gives information about the viability of the fetus, and in later stages of pregnancy gives information about the health of the fetus. At the early stage of pregnancy, it is possible to see signs of the fetal heart functioning through the use of ultrasound, while Doppler devices can assess the heart rate during standard appointments. During high-risk pregnancies, cardiotocography is used in order to evaluate the heart rate and its relation to uterine activity. [22]

4.5.3 Doppler and Other Fetal Assessments

Doppler ultrasound is a unique method that is used for examining blood circulation in some of the fetal and maternal blood vessel systems. This method is of big importance in pregnancies where there is a chance of fetal growth failure, placental inefficacy, and maternal or fetal complications. [23] There is a possibility of getting important data about the placental and fetal circulation from the assessment of blood circulation in umbilical artery and other fetal or maternal vessels if it is necessary. Doppler results can help physicians understand whether further examination is needed. Other fetal evaluations may include checking the levels of amniotic water, movements, tone, breathing movements, and heart beats of the unborn child. Biophysical profile refers to the process of using some ultrasound parameters along with fetal

heart beats assessment to estimate the health of the fetus in the pregnancy when some degree of monitoring is necessary. In some cases, non-stress test and CTG can be used if there is an indication, especially in risky pregnancies. The choice should be made according to the situation and the tests should not be used instead of routine prenatal assessment. [24]




Fig 1: Ultrasonography and Fetal Monitoring During Pregnancy

4.6 Maternal Health Assessment

As mentioned in (Table 3) the assessment of maternal health is an important part of prenatal care and encompasses the ongoing examination of both the mother's physical condition and her dietary status throughout the course of the pregnancy. The pregnancy process involves normal physiological alterations affecting various systems of the human organism, which means that continuous assessment is a means of differentiating between medical phenomena anticipated, and signs that may require additional inspection and testing. During each treatment session the healthcare provider goes through the patient's medical and obstetric history, and examines existing complaints, clinical records and risk indicators. The presence of a thorough assessment of the maternal condition allows healthcare practitioners to detect the occurrence of complications and perform timely investigations and treatments. [25]

4.6.1 Vital Signs and Physical Assessment

Frequent monitoring of vital signs of the mother provides valuable information regarding the health of the mother and may facilitate early detection of complications that can happen during pregnancy. Monitoring blood pressure has to be performed during tours to antenatal clinics, as high blood pressure may indicate pregnancy hypertension. In addition, it will be useful to monitor other parameters like pulse rate, respiratory rate, temperature, and weight, depending

on the clinical situation. Physical examination may include checking for pallor, signs of edema, or some signs of infection and other important clinical signs. Additionally, abdominal examination is done during pregnancy in order to monitor the uterine size, fetal growth, fetal lie and presentation, and heart activity. [26]

4.6.2 Nutritional Status

The evaluation of women's nutrition conditions is relevant for its connections with women's health, placentation, fetal development, labor, and lactation. Assessment of nutrition requires measuring BMI before or in early pregnancy, weight changes during pregnancy, dietary habits, and critical health parameters reflecting possible deficiency in nutrition factors. The most crucial nutrients to be monitored include calories, proteins, iron, folates, calcium, iodine, vitamin D, among many others. Both low and high weight gain during pregnancy leads to adverse results and should be corrected through dietary and lifestyle changes. Women suffering from insufficient consumption of nutrients, anemia, malnutrition, obesity, and other issues with nutrition may need additional testing and nutritional advice. [27]

4.6.3 Screening for Anemia, Diabetes and Hypertension

Screening for anemia, diabetes, and hypertensive disorders is crucial in assessing maternal health since those disorders are quite prevalent and can have a serious impact on the outcome of pregnancy if left undiagnosed or improperly treated. [28] Anemia is usually diagnosed through measuring hemoglobin and pursuing additional investigations in case the cause is uncertain. Screening for gestational diabetes occurs at certain intervals, depending on the day of pregnancy, and may include performing an oral glucose tolerance test or other tests recommended by specialists. Women with any risk factors may have to undergo the screening procedure earlier rather than at the standard date. Stating the facts concerning blood pressure in expectation of other necessary tests for women with high blood pressure can add to the screening process for preeclampsia. [29]

Table 3: Maternal Health Assessment during Pregnancy

S. No.	Assessment Area	Key Components
1	**Physical Health Assessment**	Ongoing assessment of the mother's physical condition throughout pregnancy
2	**Dietary/Nutritional Assessment**	Assessment of dietary status and nutritional requirements during pregnancy
3	**Medical and Obstetric History**	Review of the woman's medical and obstetric history during antenatal visits
4	**Clinical Assessment**	Evaluation of existing complaints, clinical records and pregnancy-related risk indicators

5	**Detection of Complications**	Identification of complications and abnormalities requiring further investigation
6	**Timely Investigation and Treatment**	Appropriate investigations and treatment when abnormalities or complications are identified

4.7 Vaccination during Pregnancy

As shown in (Fig 2) Vaccination during pregnancy is considered an essential part of preventive antenatal care as it helps to keep the mother safe from vaccine-preventable diseases and, in some cases, protect the fetus and newborn through the transfer of protective antibodies from their mother. Pregnancy can modify the immune response and increase the chance of developing complications related to specific infections. For this reason, it is essential to assess the vaccination status early during pregnancy and provide recommendations and necessary vaccines according to state immunization guidelines, mother's risk factors, pregnant status, and epidemiology of the disease. Moreover, healthcare specialists should review the vaccination history of the mother and indications prior to vaccination. [30]

4.7.1 Recommended Vaccines

Vacination can safely take place during pregnancy and some vaccines can be administered. It is advisable that individuals receive vaccination against influenza during pregnancy since pregnant women are suceptible to influenza-related complications. Maternal vaccination would aid in providing some protection to infants early in their lives. Maternal vaccination against tetanus, diphtheria and pertussis (Tdap) is also advisable since it would prevent pregnant women from suffering from these diseases and passively confer immunity to newborns. Vaccination against diphtheria and tetanus may be a part of the regular maternal vaccination program in a certain country. Furthermore, pregnant women may be advised to get vaccinated against COVID-19 depending on the regulations of a specific country on the topic and the specific situation of the woman. Hepatitis B vaccine may also be given to women with a high likelihood of infection or situation requiring immunization. Live vaccines are best avoided during pregnancy and inactive ones are mainly recommended in countries where vaccination is recommended. [31]

4.7.2 Vaccination Schedule

Vaccination timing during pregnancy varies with vaccine type, the stage of gestation, past vaccination history, and country-specific guidelines. Flu vaccines are typically given during one of the trimesters when necessary, and there is an option for seasonal flu vaccination as well. Tdap vaccine is recommended for pregnant women late in their pregnancy, and the goal in selecting the timing is to ensure maximum transfer of passive antibodies to the fetus. Vaccination schedules for tetanus and diphtheria will differ according to the patient's prior history of vaccinations and maternal immunization strategies in the local region. Ideally, a

woman's vaccination status should be checked at her first antenatal appointment to plan vaccination. The health care workers should document information regarding the vaccine used, its dose and the date of its administration. Women who miss any recommended vaccine should be assessed on a case-by-case basis. The vaccination schedule should follow the national or regional recommendations, modified according to new evidence introduced into practice, efficacy of the vaccines used, and prevalence of diseases.

4.7.3 Contraindicated Vaccines

Vaccines that are based on living agents usually are not recommended during pregnancy. Examples of such vaccines are the vaccines against measles, mumps, and rubella, the MMR vaccines, and the varicella vaccine. There are other vaccines based on living agents that may also be contraindicated unless they are carefully assessed for risks. If a live vaccine was mistakenly given before the pregnancy was established, it will usually not be a reason for terminating the pregnancy. The necessary information and follow-up are important. Vaccination decisions must take into account the information regarding previous severe allergic reactions to vaccines and their components, as well as other contraindications and precautions regarding particular vaccines. In cases when vaccinations are not recommended in pregnancy but important from the clinical point of view, they must be conducted preconceptionally or postpartum.




Fig 2: Vaccination during Pregnancy

4.8 Preventive Healthcare Measures

As shown in (Fig 3) Healthcare delivered prior to and during pregnancy is meant to help prevent diseases and other complications among both mother and child, while the correct diet, hygiene measures, physical activity, as well as healthy habits are important as well. It is known that pregnancy increases the body's need for nutrients which is accompanied by various physiologic changes with regard to the risk of infections and diseases. Thus, preventive measures must be incorporated into any routine prenatal activity starting from the early stages of pregnancy. The measures shall be tailored to fit the health and nutrition of the mother, gestational age, and potential health risks during pregnancy.

4.8.1 Nutritional Supplementation

Adequate nutrition and proper supplementation play a vital role in supporting the wellness of mothers, the growth of placentas, the innateness of embryos, readiness for childbirth, and breastfeeding. Pregnant women should get advice on a balanced diet providing sufficient energy and protein along with fruits, vegetables, whole grains, and other products rich in various nutrients. Some important nutrients, including iron, folic acid, calcium, iodine, vitamin D and some other micronutrients might need supplementation depending on individual needs and national recommendations. Iron and folic acid supplementation is especially critical in order to avoid and manage nutritional anemia and ensure proper embryo development. In particular folic acid is crucial in order to avoid neural tube defects during periconception period and early pregnancy. In case the diet does not provide sufficient amount of calcium or in case of the recommendations some local guidelines recommend calcium supplements. [32]

4.8.2 Infection Prevention

Pregnancy necessitates that particular emphasis be placed on the prevention of infections. This is due to the fact that several maternal infections can lead to serious complications for the mother or the fetus. The prevention measures should involve vaccination and hygiene practices, safe food and water handling, as well as avoiding areas with known risk factors. The pregnant women should be encouraged to take cooked food, avoid unpasteurised products, wash fruits and vegetables sufficiently, and handle food safely. It is also obligatory to take precautions when dealing with sick people who have had a contagious infection. Testing for certain infections such as HIV, hepatitis B, and syphilis has to be carried out following suggested antenatal guidelines with the provision of timely medical assistance. Safe sexual practices and barrier protection methods should also be recommended whenever sexually transmitted diseases can happen. [33]

4.8.3 Hygiene and Oral Health

Good personal hygiene practices during pregnancy can reduce infection risk and produce a sense of wellbeing and comfort. Taking regular baths, washing hands properly, practicing

genital hygiene, wearing clean clothes, and observing menstrual hygiene are to be encouraged. Advice about dental hygiene is also needed by pregnant women, since hormonal changes may put them at increased risk of gum disease or other dental problems. Through regular brushing of teeth with fluoride paste, effective flossing, proper diet, and a visit to a dentist, oral health can be maintained.

4.8.4 Physical Activity and Healthy Lifestyle

It has been established that regular exercise is important for many women experiencing uncomplicated pregnancies and that it can improve fitness, mental health, gestational weight gain, and pregnancy health in general. Among the activities that can be recommended are walking, swimming, and exercising in ways suitable for the situation, which needs to be discussed with the doctor. Activities with high-risk rates of injuries should be avoided and exercising should be adapted according to restrictions imposed by pregnancy and medical recommendations. It is also important for the mother to sleep enough and relax adequately during the whole period of pregnancy. A healthy lifestyle during pregnancy should include a rational diet, staying physically active, having enough sleep, being on time for prenatal visits, and taking all medications prescribed. A pregnant woman should not smoke or drink alcohol, as well as use any harmful substances that can negatively impact maternal health. Moreover, self-medication and taking herbs or other alternatives without prior consultation with the doctor should be avoided. Thus, to keep mother healthy and help her give birth to a healthy child, it is important to follow the principles of health living and regularly visit the doctor.

4.9 Patient Education and Counselling

As shown in (Fig 4) Patient education and counselling are key elements of prenatal care due to their role in giving clarity regarding the physiological and emotional changes during pregnancy. Counselling should be continuous and tailored to the stage of pregnancy, woman's health condition, personal requirements and risk factors. Healthcare providers should deliver specific and clear answers related to diet, medications, preventive measures, alarming symptoms, delivery preparation, lactation, psychological condition, and postpartum care. With effective counselling, patients might have better compliance with prenatal care, build healthier lifestyles, and identify acute health issues in the early stages. [34]

4.9.1 Nutrition and Medication Counselling

The importance of a well-balanced and diversified approach to nutrition must be highlighted during nutrition consultation. It is important to inform the pregnant women about the intake of healthy food items and follow the recommendations for the consumption of iron, folic acid, calcium, iodine, vitamin D, and other supplements when necessary. Food safety, adequate fluid intake, proper weight gain during pregnancy in addition to the rejection of potentially harmful products must also be discussed during consultation. Specific recommendations for making

dietary plans must be based on individual patterns of diet, health conditions, and spiritual inclinations. Another important component of the counseling process is the explanation of drug-related issues. Women need to be aware of the fact that they may take prescribed or over-the-counter medicines, food supplements, or herbal preparations. It is vital to warn women about the importance of taking medication only after consulting with a doctor. Health professionals need to make sure that medications are not dangerous during pregnancy and explain when, how long and in what dosage the medications must be taken. [35]




Fig 3: Preventive Healthcare measures during Pregnancy

4.9.2 Warning Signs

Education for the pregnant should include information about symptoms signaling a significant maternal or fetal problem and the need for immediate medical assistance. Symptoms that necessitate quick evaluation by doctors include any vaginal bleeding, fluid leakage, severe abdominal pain, severe headache, visual disturbances, sudden swelling of the face or hands, shortness of breath, chest pain, high fever, seizures, severe vomiting, or drastically reduced or absent fetal movements if they were present before. Women must be educated to hurry to seek medical aid in the occurrence of such a symptom. They will need the data about the nearest hospitals to turn to and the time of hospitalization.

4.9.3 Birth Preparedness

The process of preparing for childbirth involves making arrangements beforehand for both labor and unexpected emergencies. For counseling to be comprehensive, it should focus on aspects such as deciding on the place where the baby will be delivered, choosing a suitable health service provider, planning travel to the place of delivery, arranging for someone to help during the delivery process, preparing necessary paperwork and equipment and taking care of issues related to finances, if needed. Expectant mothers also need to have knowledge of labor as well as the timing for contacting health services if necessary. In cases of high-risk pregnancy, women should develop a personalized plan with healthcare professionals to ensure they will deliver their babies at a facility with sufficient maternal and neonatal care capabilities. Counseling should also include the aspects of breastfeeding and proper newborn care.

4.9.4 Mental Health and Family Support

The hormonal changes associated with pregnancy can cause a wide range of emotions, anxiety, tension, and worries regarding childbirth, health, responsibility towards the family, and social circumstances. Therefore, prenatal counseling has to consider the emotional and psychological well-being of the mother. Healthcare providers need to motivate future mothers to identify constant sadness, anxiety, sleep problems, lack of interest, or other emotional problems. The women who are suffering from severe psychological discomfort need proper evaluation and referral for psychological help when needed. Family and partner support can provide a positive contribution to a woman's well-being and compliance with prenatal care. The family can help in terms of proper nutrition and taking medicines and supplements, ensuring transportation to appointments and home chores, and preparing for childbirth. When the woman agrees to this approach, the inclusion of family into adequate counseling ensures the realization of what the future mother may need and a supportive atmosphere in the family.

4.10 Role of Healthcare Professionals

The healthcare professionals are vital in ensuring the provision of safe, effective, and continuous prenatal healthcare services throughout the duration of pregnancy. Similarly, obstetricians, doctors, nurses, midwives, pharmacists, nutritionists, and other members of the healthcare team also support maternal and fetal health through their regular assessments, examinations, preventive care, treatment, education, and counseling. Healthcare professionals monitor maternal parameters such as vital signs, nutrition, weight gain, fetal growth, and laboratory values so as to identify various risk factors and symptoms of complications in pregnancy like anemia, gestational diabetes, hypertension, and infections.

Medical professionals should also inform expectant mothers on the topics of nutrition, safe and effective use of drugs, exercise, hygiene, immunization, signs of illness during pregnancy, preparedness for delivery, and caring for the infant after birth. They should motivate women to

keep their doctor appointments and use prescription drugs as recommended. Also, the healthcare workers must resolve any questions and misunderstandings in a respectful and easy way. Pharmacists can help by checking the medication supplied to the pregnant women and highlighting potential problems related to drugs. Nutritionists can give personalized advice on diet based on the previous diet of the pregnant women. Nurses and midwives continue monitoring, counseling, providing emotional support, and coordinating care.




Fig 4: Components of Patient Education and Counseling during Pregnancy

Detection of the problems and timely referrals is the main duty of the healthcare personnel working with women who have complicated pregnancies. Collaboration between different specialists guarantees continuity of provision of services and proper care of women during pregnancy.

4.11 References

1. Arsenault C, Mfeka-Nkabinde NG, Chaudhry M, Jarhyan P, Taddele T, Mugenya I, Sabwa S, Wright K, Amboko B, Baensch L, Molla Wondim G. Antenatal care quality and detection of risk among pregnant women: An observational study in Ethiopia, India, Kenya, and South Africa. PLoS medicine. 2024 Aug 27;21(8):e1004446.
2. McCauley H, Lowe K, Furtado N, Mangiaterra V, van den Broek N. What are the essential components of antenatal care? A systematic review of the literature and development of

signal functions to guide monitoring and evaluation. BJOG: An International Journal of Obstetrics & Gynaecology. 2022 May;129(6):855-67.

3. Redi T, Seid O, Bazie GW, Amsalu ET, Cherie N, Yalew M. Timely initiation of antenatal care and associated factors among pregnant women attending antenatal care in Southwest Ethiopia. PloS one. 2022 Aug 18;17(8):e0273152.
4. Nam JY, Oh SS, Park EC. The association between adequate prenatal care and severe maternal morbidity among teenage pregnancies: a population-based cohort study. Frontiers in Public Health. 2022 May 31;10:782143.
5. Goncalves AS, Ferreira IM, Pestana-Santos M, McCourt C, Prata AP. Antenatal care policy in high-income countries with a universal health system: A scoping review. Sexual & reproductive healthcare. 2022 Jun 1;32:100717.
6. Balk EM, Konnyu KJ, Cao W, Reddy Bhuma M, Danilack VA, Adam GP, Matteson KA, Peahl AF. Schedule of visits and televisits for routine antenatal care: a systematic review.
7. Menon G, CP S, PK A. Yoga in Pregnancy: Trimester-Wise Evidence based Yoga Protocol for Pregnant women. Journal of Chemical Health Risks. 2025 Dec 8;15(5):3404.
8. Pi X, Wang J, Chu L, Zhang G, Zhang W. Prediction of high-risk pregnancy based on machine learning algorithms. Scientific Reports. 2025 May 4;15(1):15561.
9. Jahan N, Akther J. Machine Learning Approaches for Early Prediction of High-Risk Pregnancy in Underserved US Populations. American Journal of Interdisciplinary Studies. 2024 Dec 28;5(04):268-309.
10. Alim A, Imtiaz MH. Wearable sensors for the monitoring of maternal health—a systematic review. Sensors. 2023 Feb 22;23(5):2411.
11. Bohn MK, Adeli K. Physiological and metabolic adaptations in pregnancy: importance of trimester-specific reference intervals to investigate maternal health and complications. Critical Reviews in Clinical Laboratory Sciences. 2022 Feb 17;59(2):76-92.
12. David AL, Spencer RN. Clinical assessment of fetal well-being and fetal safety indicators. The Journal of Clinical Pharmacology. 2022 Sep;62:S67-78.
13. Jain S, Acharya N. Fetal wellbeing monitoring: a review article. Cureus. 2022 Sep 11;14(9).
14. Allanazarov I, Rahmonova U, Abdumannonova K. PROBLEMS OF PREVENTION AND EARLY DIAGNOSIS OF COMPLICATIONS DURING PREGNANCY. INTEGRATION OF EDUCATION AND SCIENCE: GLOBAL CHALLENGES AND SOLUTIONS. 2025 Nov 14;1(2):48-52.

15. Ertmann RK, Nicolaisdottir DR, Kragstrup J, Overbeck G, Kriegbaum M, Siersma V. The predictive value of common symptoms in early pregnancy for complications later in pregnancy and at birth. Acta Obstetricia et Gynecologica Scandinavica. 2023 Jan;102(1):33-42.

16. Abbassi-Ghanavati M, Greer LG, Cunningham FG. Pregnancy and laboratory studies: a reference table for clinicians. Obstetrics & Gynecology. 2009 Dec 1;114(6):1326-31.

17. Patel P, Balanchivadze N. Hematologic findings in pregnancy: a guide for the internist. Cureus. 2021 May 21;13(5).

18. Bi BC, Yang HY, Su JY, Deng L. Analysis of pregnancy and neonatal outcomes in 100 pregnant women with Rh-negative blood type. BMC Pregnancy and Childbirth. 2024 Dec 18;24(1):815.

19. Ahmad S, Tabassum H, Mobeen I, Mirza A, Hurain A, Zulfiqar H, Noor M. Clinical Evaluation of Urinalysis and Glucose Levels in Pregnancy: Implications for Maternal Health. Journal of Pharma and Biomedics. 2026 Mar 31;4(2):65-71.

20. Wu S, Wang J, Guo Q, Lan H, Sun Y, Ren M, Liu Y, Wang P, Wang L, Su R, Zhang J. Prevalence of human immunodeficiency virus, syphilis, and hepatitis B and C virus infections in pregnant women: a systematic review and meta-analysis. Clinical microbiology and infection. 2023 Aug 1;29(8):1000-7.

21. Khalil A, Sotiriadis A, D'Antonio F, Da Silva Costa F, Odibo A, Prefumo F, Papageorghiou AT, Salomon LJ. ISUOG Practice Guidelines: performance of third-trimester obstetric ultrasound scan. Ultrasound in Obstetrics & Gynecology. 2024;63(1):131-47.

22. Ahmed MR, Newby S, Potluri P, Mirihanage W, Fernando A. Emerging paradigms in fetal heart rate monitoring: evaluating the efficacy and application of innovative textile-based wearables. Sensors. 2024 Sep 19;24(18):6066.

23. Sá MI, Illa M, Guedes-Martins L. Doppler assessment of the fetal brain circulation. Diagnostics. 2026 Jan 9;16(2):214.

24. Ochoa JH, Cafici D. Fetal Doppler assessment in pregnancy. Best Practice & Research Clinical Obstetrics & Gynaecology. 2025 Jun 1;100:102594.

25. Khadidos AO, Saleem F, Selvarajan S, Ullah Z, Khadidos AO. Ensemble machine learning framework for predicting maternal health risk during pregnancy. Scientific Reports. 2024 Sep 14;14(1):21483.

26. Draves J, TEKİNER H, Yale ES, Mazza JJ, Yale SH. A comprehensive assessment of the eight vital signs. Eurobiotech Journal. 2022;6(3).

27. Mohamed HJ, Loy SL, Mitra AK, Kaur S, Teoh AN, Rahman SH, Amarra MS. Maternal diet, nutritional status and infant birth weight in Malaysia: a scoping review. BMC pregnancy and childbirth. 2022 Apr 6;22(1):294.

28. Stanley AY, Wallace JB, Hernandez AM, Spell JL. Anemia in pregnancy: screening and clinical management strategies. MCN: The American Journal of Maternal/Child Nursing. 2022 Jan 1;47(1):25-32.

29. Chauhan BG, Chokhandre PK, Kulaste BS, Sivanandan V. Burden of anaemia, hypertension and diabetes among pregnant women in India. Journal of Biosocial Science. 2023 Sep;55(5):931-46.

30. De Weerdt L, Herzog SA, Van Damme P, Maertens K. Timing of pertussis vaccination during pregnancy: evidence and implementation–a systematic review. Vaccine. 2024 Aug 30;42(21):126152.

31. Licata F, Romeo M, Riillo C, Di Gennaro G, Bianco A. Acceptance of recommended vaccinations during pregnancy: a cross-sectional study in Southern Italy. Frontiers in Public Health. 2023 May 12;11:1132751.

32. Liu E, Wang D, Darling AM, Perumal N, Wang M, Ahmed T, Christian P, Dewey KG, Kac G, Kennedy S, Subramoney V. Effects of prenatal nutritional supplements on gestational weight gain in low-and middle-income countries: a meta-analysis of individual participant data. The American Journal of Clinical Nutrition. 2022 Dec;116(6):1864-76.

33. Yasin R, Jiang L, Das JK, Bhutta ZA. Interventions to prevent and manage infections in pregnancy. Neonatology. 2025 Feb 28;122(Suppl. 1):32-41.

34. Al-Mutawtah M, Campbell E, Kubis HP, Erjavec M. Women's experiences of social support during pregnancy: a qualitative systematic review. BMC pregnancy and childbirth. 2023 Nov 10;23(1):782.

35. Dewidar O, John J, Baqar A, Madani MT, Saad A, Riddle A, Ota E, Kung'u JK, Arabi M, Raut MK, Klobodu SS. Effectiveness of nutrition counseling for pregnant women in low-and middle-income countries to improve maternal and infant behavioral, nutritional, and health outcomes: A systematic review. Campbell systematic reviews. 2023 Dec;19(4):e1361.